

# Hydrogen isotopic ratios of long-chain diols reflect salinity

## Journal Article

### Author(s):

Lattaud, Julie  Erdem, Zeynep; Weiss, Gabriella M.; Rush, Darci; Balzano, Sergio; Chivall, David; van der Meer, Marcel T. J.; Hopmans, Ellen C.; Sinninghe Damsté, Jaap S.; Schouten, Stefan

### Publication date:

2019-11

### Permanent link:

<https://doi.org/10.3929/ethz-b-000386416>

### Rights / license:

Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International

### Originally published in:

Organic Geochemistry 137, <https://doi.org/10.1016/j.orggeochem.2019.103904>



## Note

## Hydrogen isotopic ratios of long-chain diols reflect salinity

Julie Lattaud<sup>a,\*</sup>, Zeynep Erdem<sup>a</sup>, Gabriella M. Weiss<sup>a</sup>, Darci Rush<sup>a</sup>, Sergio Balzano<sup>a,1</sup>, David Chivall<sup>a,2</sup>, Marcel T.J. van der Meer<sup>a</sup>, Ellen C. Hopmans<sup>a</sup>, Jaap S. Sinninghe Damsté<sup>a,b</sup>, Stefan Schouten<sup>a,b</sup>

<sup>a</sup> NIOZ Royal Netherlands Institute for Sea Research, Department of Marine Microbiology and Biogeochemistry, and Utrecht University, the Netherlands

<sup>b</sup> Utrecht University, Department of Earth Sciences, Faculty of Geosciences, Princetonlaan 8A, 3584 CB Utrecht, the Netherlands

## ARTICLE INFO

## Article history:

Received 20 June 2019

Received in revised form 30 July 2019

Accepted 2 August 2019

Available online 5 August 2019

## Keywords:

Hydrogen isotopes

Long-chain diols

Salinity

Preparative HPLC

## ABSTRACT

Long-chain diols (LCDs) are ubiquitous lipids produced by freshwater and marine algae. A combination of semi-preparative high performance liquid chromatography and gas chromatography isotope ratio monitoring mass spectrometry, allowed the measurement of  $\delta^2\text{H}$  of individual LCDs from cultures, which indicated a correlation with the hydrogen isotope composition of the growth water and a species-specific effect. Results from environmental samples along a salinity gradient indicated the potential of  $\delta^2\text{H}$  ratios of LCDs to trace the hydrogen isotopic composition of water and sea surface salinity.

© 2019 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Hydrogen isotopes ratios of lipids from photosynthetic organisms have been shown to trace the hydrogen isotopic composition of the water ( $\delta^2\text{H}_{\text{H}_2\text{O}}$ ) as well as salinity in cultures (e.g., Schouten et al., 2006; Zhang and Sachs, 2007; Zhang et al., 2009; Sachs et al., 2016; Weiss et al., 2017). This principle has been used in particular with  $\delta^2\text{H}$  ratios of long-chain alkenones as a proxy for  $\delta^2\text{H}_{\text{H}_2\text{O}}$  and sea surface salinity (van der Meer et al., 2007; Weiss et al., 2019 and references therein). However, there are few other marine biomarkers used for this purpose.

Long-chain diols (LCDs) are ubiquitous marine compounds possibly produced by eustigmatophytes (Volkman et al., 1992, 1999; Villanueva et al., 2014). The  $\text{C}_{28}$  and  $\text{C}_{30}$  1,13- and the  $\text{C}_{30}$  1,15-diols are the major marine LCDs and have been used in a sea surface temperature proxy, the Long-chain Diol Index (Rampen et al., 2012). Furthermore, the fractional abundance of the  $\text{C}_{32:0}$  1,15-diol has been used to trace riverine input in marginal seas as it is mainly produced in river systems by freshwater eustigmatophytes (Lattaud et al., 2017), and is in low abundance in open marine environments (Rampen et al., 2012). Thus,  $\delta^2\text{H}$  ratios of

LCDs might be potential tools to reconstruct the  $\delta^2\text{H}_{\text{H}_2\text{O}}$  of sea water as well as sea surface salinity and  $\delta^2\text{H}_{\text{H}_2\text{O}}$  of river water, respectively. However, LCD isomers are not separated by traditional gas chromatography (GC) (cf. Versteegh et al., 1997) so semi-preparative liquid chromatography (prep-LC) is necessary to obtain  $\delta^2\text{H}$  ratios of individual LCDs (De Bar et al., 2016).

Here we used a prep-LC method adapted from de Bar et al. (2016) to isolate individual LCDs and measured their  $\delta^2\text{H}_{\text{LCDs}}$  in cultures and marine surface sediments from an environmental salinity gradient.

## 2. Materials and methods

## 2.1. Cultures and surface sediments

Three strains of *Nannochloropsis* (*Nannochloropsis gaditana* strain CCMP526, *Nannochloropsis oceanica* strain CCMP1779, and *Nannochloropsis oculata* strain CCMP2195) were grown in triplicate with a light intensity of  $280 \mu\text{mol m}^{-2} \text{s}^{-1}$  and salinity of  $37.5 \text{ g kg}^{-1}$  following methods described in Balzano et al. (2017). *N. oculata* was also grown at five salinities (22.5, 27.8, 30.1, 32.7 and  $34.5 \text{ g kg}^{-1}$ ) at  $15^\circ\text{C}$  and a light intensity of  $60 \mu\text{mol m}^{-2} \text{s}^{-1}$ . Cell counts were obtained by flow cytometry (BD Accuri™ C6). The cells were harvested in exponential or, for the salinity experiment, stationary growth phase by filtration through  $0.7 \mu\text{m}$  GF/F filters (Whatman, Maidstone, UK).

Six surface sediments (Fig. 1) were collected in the northern Gulf of Mexico in March 2018 using a multi-corer (R/V *Pelagia*

\* Corresponding author at: Department of Earth Sciences, ETH Zurich, Sonneggstrasse 5, 8006 Zurich, Switzerland.

E-mail address: [jlattaud@ethz.ch](mailto:jlattaud@ethz.ch) (J. Lattaud).

<sup>1</sup> Present address: Stazione Zoologica Anton Dohrn, Istituto Nazionale di Biologia, Ecologia e Biotecnologie Marine, Villa Comunale, 80121 Naples, Italy.

<sup>2</sup> Present address: Oxford Radiocarbon Accelerator Unit, School of Archaeology, University of Oxford, OX1 3TG, UK.




Fig. 1. (a) Location of the surface sediments and sea surface salinity (Antonov et al., 2010) and (b)  $\delta^2\text{H}$  ratios of the lipids.

cruise 64PE434). The annual mean salinity and annual mean temperature varies from 33.4 to 35.2 g kg<sup>-1</sup> and from 24.5 to 25.4 °C, respectively (Locarnini et al., 2009; Antonov et al., 2010).

ples were analyzed in triplicate when possible. The  $\delta^2\text{H}$  ratios of the LCDs were corrected for BSTFA (surface sediments) or acetylation (for culture experiments) as followed:

$$\delta^2\text{H}_{\text{LCD}} = \frac{(\text{nH}_{\text{LCD}} + 2 \times \text{nH}_{\text{derivative}}) \times \delta^2\text{H}_{\text{LCD measured}} - 2 \times \text{nH}_{\text{derivative}} \times \delta^2\text{H}_{\text{derivative}}}{\text{nH}_{\text{LCD}}} \quad (1)$$

## 2.2. Lipid extraction and separation

All surface sediments were freeze-dried and extracted using Accelerated Solvent Extraction following Lattaud et al. (2017). The filters from *Nannochloropsis* cultures were extracted following Lattaud et al. (2018) using base and acid hydrolysis. An internal standard (C<sub>22</sub> 7,16-diol) was added to all total lipid extracts and all extracts were subsequently separated into three fractions (apolar, ketone and polar) on an Al<sub>2</sub>O<sub>3</sub> column as described in Lattaud et al. (2018). For the quantification of the LCDs the polar fractions were analyzed by GC–MS as described in Lattaud et al. (2018). For the salinity experiment, saturated LCDs were isolated from the polar fraction by elution through an Ag<sup>+</sup> impregnated Si column. The polar fractions were then acetylated with acetic anhydride ( $\delta^2\text{H} = -125.8 \pm 3.3\text{‰}$ ) following Das and Chakraborty (2011).

## 2.3. Isolation of long-chain diol isomers

Prior to stable isotope analysis, the LCDs were isolated from the polar fraction following a method from De Bar et al. (2016) using semi-preparative normal phase High Performance Liquid Chromatography (HPLC). An Agilent 1260 infinity HPLC equipped with a thermostatted autoinjector, and a Foxy R1 fraction collector with a Sunfire silica column (Waters, 250 mm × 10 mm; 5 μm) was used and 20 s fraction were collected. Fractions were combined in three pools: with pool 1 (33.4–34.4 min) containing 100% of C<sub>32:0</sub> 1,15-diol, pool 2 (35–35.2 min) containing 95% of C<sub>30</sub> 1,15-diol and pool 3 (35.4–37.4 min) containing 100% of C<sub>28</sub> 1,13 and <90% of C<sub>28</sub> 1,14-diols. Hydrogen isotope fractionation during HPLC isolation was tested using a pure LCD standard with known hydrogen isotope composition (C<sub>28</sub> 1,13-diol;  $\delta^2\text{H} = -142.6 \pm 0.6\text{‰}$ ). Prior to isotopic measurements, the purified LCDs were silylated with BSTFA with a known hydrogen isotopic composition ( $\delta^2\text{H} = -212 \pm 1\text{‰}$ ).

## 2.4. $\delta^2\text{H}$ measurements

Isolated LCDs from the polar fraction and long-chain alkenones from the ketone fraction were analyzed using gas chromatography isotope ratio monitoring mass spectrometry (GC–irMS) using a CPSil5 GC column (Agilent, 25 m × 0.32 mm × 0.4 μm). The sam-

With  $\text{nH}_{\text{LCD}}$  = number of hydrogen atoms of LCD, and the derivative being either BSTFA or acetyl anhydride.  $\text{nH}_{\text{BSTFA}} = 9$  and  $\text{nH}_{\text{acetyl}} = 3$ ,  $\delta^2\text{H}_{\text{BSTFA}} = -212\text{‰}$  and  $\delta^2\text{H}_{\text{acetyl}} = -125.8\text{‰}$ .  $\delta^2\text{H}_{\text{BSTFA}}$  has been determined by measuring the  $\delta^2\text{H}$  of a BSTFA-derivatized phthalic acid with known isotopic ratio ( $-81.9 \pm 1.2\text{‰}$ ).

## 3. Results and discussion

### 3.1. Hydrogen isotopic composition of LCDs in cultures

A C<sub>28</sub> 1,13-diol standard was collected over three separate fractions using prep-LC containing 17, 59 or 21% of the original compound, respectively. A large isotope effect was noted ( $\delta^2\text{H} = -190.5 \pm 1.9\text{‰}$ ,  $-152.1 \pm 0.9\text{‰}$ ,  $-101.7 \pm 1.0\text{‰}$ , respectively) as observed previously for other compounds (Atwood and Sachs, 2012). Thus, it is necessary to ensure that 95% of the targeted peak is recombined to avoid any chromatographic fractionation effect ( $\leq 5\text{‰}$ ).

We analyzed three different *Nannochloropsis* species to assess whether there are differences in  $\delta^2\text{H}_{\text{LCD}}$  between species. The cultures contained mainly the C<sub>32:1</sub> 1,15- and C<sub>32:0</sub> 1,15-diols as reported previously (Volkman et al., 1992; Balzano et al., 2017), and no C<sub>30</sub> 1,15-diol. In *N. oculata* and *N. gaditana*, the C<sub>32:0</sub> 1,15-diol was more abundant than the C<sub>32:1</sub> 1,15-diol, whereas the latter was the major LCD in *N. oceanica*. The  $\delta^2\text{H}$  ratios of the C<sub>32:1</sub> 1,15- and the C<sub>32:0</sub> 1,15-diols did not differ significantly between *N. oceanica* and *N. oculata* (Table 1). In contrast, the  $\delta^2\text{H}$  of the C<sub>32:1</sub> 1,15-diol was more depleted ( $-135 \pm 6\text{‰}$ ) than the C<sub>32:0</sub> 1,15-diol ( $-129 \pm 2\text{‰}$ , Table 1) in *N. gaditana*, and both values were depleted considerably relative to the corresponding values in the other two species, suggesting that there is a species effect on the hydrogen isotopic fractionation of LCDs. Salinity experiments with *N. oculata* showed that the  $\delta^2\text{H}$  ratio of the C<sub>32:0</sub> 1,15-diol is more enriched at higher salinity, but that fractionation is relatively constant with slightly lower values at the highest salinity (Table 2). These observations indicate that  $\delta^2\text{H}_{\text{LCD}}$  might reflect the  $\delta^2\text{H}$  of growth medium, and potentially salinity, in which LCD-producers grow. The  $\delta^2\text{H}$  ratios of *N. oculata* LCDs in the salinity experiment are enriched compared to the initial culture grown at 37.5 g kg<sup>-1</sup> (Table 1), suggesting other factor such as light and growth phase

**Table 1**

$\delta^2\text{H}$  ratios of *Nannochloropsis* spp. cultures with standard deviation, and fractionation values between the lipids and the medium indicated as  $\alpha$ .

Names	CCMP 526 ( <i>N. gaditana</i> )	CCMP 1779 ( <i>N. oceanica</i> )	CCMP 2195 ( <i>N. oculata</i> )
$\delta^2\text{H}_{\text{C}_{32:0} \ 1,15} \ (‰)$	$-129 \pm 2$	$-118 \pm 10$	$-116 \pm 4$
$\delta^2\text{H}_{\text{C}_{32:1} \ 1,15} \ (‰)$	$-135 \pm 6$	$-114 \pm 2$	$-116 \pm 3$
$\delta^2\text{H}_{\text{medium}} \ (‰)$	8.9	8.7	8.9
$\alpha$	0.863	0.874	0.876

**Table 2**

$\delta^2\text{H}$  ratios of *N. oculata* cultures with standard deviation (samples without a value were only measured once due to low concentration), and fractionation values between the lipids and the medium indicated as  $\alpha$ . Salinity and  $\delta^2\text{H}_{\text{medium}}$  are the mean of measurements made at the start of the culture and at harvesting.

Salinity (g kg <sup>-1</sup> )	22.5	27.8	30.1	32.7	34.5
$\delta^2\text{H}_{\text{C}_{32:0} \ 1,15} \ (‰)$	$-118 \pm 11$	$-116 \pm 3$	-105	$-105 \pm 6$	-95
$\delta^2\text{H}_{\text{medium}} \ (‰)$	$-13.7 \pm 5.4$	$-8.1 \pm 6.8$	$-1.8 \pm 3.5$	$+0.8 \pm 6.7$	$+4.0 \pm 3.4$
$\alpha$	0.894	0.891	0.897	0.894	0.901

likely impact  $\delta^2\text{H}_{\text{LCD}}$  (Wolhowe et al., 2009; Zhang et al., 2009; Chivall et al., 2014; van der Meer et al., 2015).

### 3.2. Hydrogen isotopic composition of LCDs in marine surface sediments

The  $\delta^2\text{H}$  ratio of the  $\text{C}_{32:0}$  1,15-diol in surface sediments from the Gulf of Mexico were similar to each other ( $-260 \pm 5‰$ ,  $n = 4$ , Fig. 1) whereas the  $\delta^2\text{H}$  ratios of the  $\text{C}_{30}$  1,15 diol was more enriched and increased with salinity (from  $-207‰$  to  $-184‰$ , Fig. 1). The large isotopic difference (up to  $70‰$ ) between the  $\text{C}_{32:0}$  1,15-diol and  $\text{C}_{30}$  1,15-diol supports the idea that the former originates mainly from river input (Lattaud et al., 2017). Indeed, the  $\text{C}_{30}$  1,15-diol concentrations were lower close to the river mouth ( $0.2$  vs  $1.0 \mu\text{g g}^{-1}$ ), in contrast to the  $\text{C}_{32:0}$  1,15-diol ( $0.1$  vs  $0.05 \mu\text{g g}^{-1}$ ). The  $^2\text{H}$ -depletion of the  $\text{C}_{32:0}$  1,15-diol likely originates from the lower  $\delta^2\text{H}_{\text{H}_2\text{O}}$  values of the Mississippi or Atchafalaya rivers ( $-31 \pm 8‰$  for the Atchafalaya River in Melville and  $-39 \pm 3‰$  for the Mississippi River in St Francisville, Lee and Veizer, 2003) rather than the marine  $\delta^2\text{H}_{\text{H}_2\text{O}} \approx 15‰$  (calculated from  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$  ratios, LeGrande and Schmidt, 2006 and the global meteoritic water line, Craig, 1961). The  $\delta^2\text{H}_{\text{C}_{30} \ 1,15}$  weakly correlated with annual mean salinity ( $r = 0.7$ ,  $p = 0.1$ ,  $n = 6$ ) and autumn mean salinity ( $r = 0.7$ ,  $p = 0.08$ ,  $n = 6$ ) in contrast to the  $\delta^2\text{H}_{\text{C}_{37}}$  ( $r = 0.49$ ,  $p = 0.4$ ,  $n = 5$ ).

## 4. Conclusions

We report, for the first time, the  $\delta^2\text{H}$  ratios of LCDs obtained from cultures of three strains of marine eustigmatophytes and environmental samples along a salinity gradient in the Gulf of Mexico. The  $\text{C}_{32:0}$  1,15-diol seems to reflect the  $\delta^2\text{H}$  ratio of the rivers/lakes where this LCD was produced while the  $\text{C}_{30}$  1,15 reflected the  $\delta^2\text{H}$  ratio and salinity of marine surface waters. The use of LCDs in conjunction with long-chain alkenones could be useful for understanding hydrological changes in the geologic record.

## Acknowledgements

We thank Co-Editor-in-Chief J. K. Volkman, Associate Editor E. Minor and an anonymous reviewer for their useful comments on this manuscript. We thank Jort Ossebaar for analytical help and the captain and crew of the R/V *Pelagia* of cruise 64PE434 (NICO

expedition, Leg 7). This research has been funded by the European Research Council (ERC) under the European Union's Seventh Framework Program (FP7/2007-2013) ERC grant agreement [339206] to S.S.

Associate Editor—Elizabeth Minor

## References

- Antonov, J.I., Seidov, D., Boyer, T.P., Locarnini, R.A., Mishonov, A.V., Garcia, H.E., 2010. In Levitus, S. (Ed.), World Ocean Atlas 2009, Volume 2: Salinity. NOAA Atlas NESDIS 69.
- Atwood, A.R., Sachs, J.P., 2012. Purification of dinosterol from complex mixtures of sedimentary lipids for hydrogen isotope analysis. *Organic Geochemistry* 48, 37–46.
- Balzano, S., Villanueva, L., de Bar, M., Sinninghe Damsté, J.S., Schouten, S., 2017. Impact of culturing conditions on the abundance and composition of long chain alkyl diols in species of the genus *Nannochloropsis*. *Organic Geochemistry* 108, 9–17.
- Chivall, D., M'Boule, D., Sinke-Schoen, D., Sinninghe Damsté, J.S., Schouten, S., van der Meer, M.T.J., 2014. The effects of growth phase and salinity on the hydrogen isotopic composition of alkenones produced by coastal haptophyte algae. *Geochimica et Cosmochimica Acta* 140, 381–390.
- Craig, H., 1961. Isotopic variations in meteoric water. *Science* 133, 1702–1703.
- de Bar, M.W., Dorhout, D.J.C., Hopmans, E.C., Rampen, S.W., Sinninghe Damsté, J.S., Schouten, S., 2016. Constraints on the application of long chain diol proxies in the Iberian Atlantic margin. *Organic Geochemistry* 101, 184–195.
- Das, R., Chakraborty, D., 2011. Silver triflate catalyzed acetylation of alcohols, thiols, phenols, and amines. *Synthesis* 2011 (10), 1621–1625.
- Lattaud, J., Kim, J.H., De Jonge, C., Zell, C., Sinninghe Damsté, J.S., Schouten, S., 2017. The  $\text{C}_{32}$  alkane-1,15-diol as a tracer for riverine input in coastal seas. *Geochimica et Cosmochimica Acta* 202, 146–158.
- Lattaud, J., Kirkels, F., Peterse, F., Freymond, C.V., Eglinton, T.I., Heftler, J., Mollenhauer, G., Balzano, B., Villanueva, L., van der Meer, M.T.J., Hopmans, E. C., Sinninghe Damsté, J.S., Schouten, S., 2018. Long-chain diols in rivers: Distribution and potential biological sources. *Biogeosciences* 15, 4147–4161.
- Lee, D., Veizer, J., 2003. Water and carbon cycles in the Mississippi River basin: potential implications for the Northern Hemisphere residual terrestrial sink. *Global Biogeochemical Cycles* 17. <https://doi.org/10.1029/2002gb001984>.
- LeGrande, A.N., Schmidt, G.A., 2006. Global gridded data set of the oxygen isotopic composition in seawater. *Geophysical Research Letters* 33, 1–5.
- Locarnini, R.A., Mishonov, A.V., Antonov, J.I., Boyer, T.P., Garcia, H.E., 2009. World Ocean Atlas 2009, Volume 1: Temperature. NOAA Atlas NESDIS 68.
- Rampen, S.W., Willmott, V., Kim, J.H., Uliana, E., Mollenhauer, G., Schefuß, E., Sinninghe Damsté, J.S., Schouten, S., 2012. Long chain 1,13- and 1,15-diols as a potential proxy for palaeotemperature reconstruction. *Geochimica et Cosmochimica Acta* 84, 204–216.
- Sachs, J.P., Maloney, A.E., Gregersen, J., Paschall, C., 2016. Effect of salinity on  $^2\text{H}/^1\text{H}$  fractionation in lipids from continuous cultures of the coccolithophorid *Emiliania huxleyi*. *Geochimica et Cosmochimica Acta* 189, 96–109.
- Schouten, S., Ossebaar, J., Schreiber, K., Kienhuis, M.V.M., Langer, G., Benthien, A., Bijma, J., 2006. The effect of temperature, salinity and growth rate on the stable hydrogen isotopic composition of long chain alkenones produced by *Emiliania huxleyi* and *Gephyrocapsa oceanica*. *Biogeosciences* 3, 113–119.
- van der Meer, M.T.J., Baas, M., Rijpsma, W.I.C., Marino, G., Rohling, E.J., Sinninghe Damsté, J.S., Schouten, S., 2007. Hydrogen isotopic compositions of long-chain alkenones record freshwater flooding of the Eastern Mediterranean at the onset of sapropel deposition. *Earth and Planetary Science Letters* 262, 594–600.
- van der Meer, M.T.J., Benthien, A., French, K.L., Epping, E., Zondervan, I., Reichart, G.-J., Bijma, J., Sinninghe Damsté, J.S., Schouten, S., 2015. Large effect of irradiance on hydrogen isotope fractionation of alkenones in *Emiliania huxleyi*. *Geochimica et Cosmochimica Acta* 160, 16–24.
- Versteegh, G.J.M., Bosch, H.-J., de Leeuw, J.W., 1997. Potential palaeoenvironmental information of  $\text{C}_{24}$  to  $\text{C}_{36}$  mid-chain diols, keto-ols and mid-chain hydroxy fatty acids: a critical review. *Science* 277, 1–13.
- Villanueva, L., Besseling, M., Rodrigo-Gámiz, M., Rampen, S.W., Verschuren, D., Sinninghe Damsté, J.S., 2014. Potential biological sources of long chain alkyl diols in a lacustrine system. *Organic Geochemistry* 68, 27–30.
- Volkman, J.K., Barrett, S.M., Blackburn, S.I., 1999. Eustigmatophyte microalgae are potential sources of  $\text{C}_{29}$  sterols,  $\text{C}_{22}$ – $\text{C}_{28}$   $n$ -alcohols and  $\text{C}_{28}$ – $\text{C}_{32}$   $n$ -alkyl diols in freshwater environments. *Organic Geochemistry* 30, 307–318.
- Volkman, J.K., Barrett, S.M., Dunstan, G.A., Jeffrey, S.W., 1992.  $\text{C}_{30}$ – $\text{C}_{32}$  alkyl diols and unsaturated alcohols in microalgae of the class Eustigmatophyceae. *Organic Geochemistry* 18, 131–138.
- Weiss, G.M., Schouten, S., Sinninghe Damsté, J.S., van der Meer, M.T.J., 2019. Constraining the application of hydrogen isotopic composition of alkenones as a salinity proxy using marine surface sediments. *Geochimica et Cosmochimica Acta* 250, 34–48.
- Weiss, M.G., Pfannerstill, Y.E., Schouten, S., Sinninghe Damsté, J.S., van der Meer, M. T.J., 2017. Effects of alkalinity and salinity at low and high light intensity on hydrogen isotope fractionation of long-chain alkenones produced by *Emiliania huxleyi*. *Biogeosciences* 14, 5693–5704.

- Wolhowe, M.D., Prah, F.G., Probert, I., Maldonado, M., 2009. Growth phase dependent hydrogen isotopic fractionation in alkenone-producing haptophytes. *Biogeochemistry* 6, 1681–1694.
- Zhang, Z., Sachs, J.P., 2007. Hydrogen isotope fractionation in freshwater algae: I. Variations among lipids and species. *Organic Geochemistry* 38, 582–608.
- Zhang, Z., Sachs, J.P., Marchetti, A., 2009. Hydrogen isotope fractionation in freshwater and marine algae: II. Temperature and nitrogen limited growth rate effects. *Organic Geochemistry* 40, 428–439.